

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/735,084	12/15/2003	Joseph C. Walsh	2003P88073US


## Response To OFFICIAL ACTION

EXAMINER	
Kriehnan, Ganapathy	
ART UNIT	PAGE NUMBER
1623	2

## AMENDMENTS TO THE CLAIMS


In the Claims, please make the following amendments:

1. (Original) A method for preparing a compound having the following formula:




wherein R is an alkoxy blocking group; P is a hydroxyl protecting group; and L is a leaving group, the method comprising the steps of:

- a. reacting a compound of the formula:




with a hydroxyl protecting group to produce a compound having the following formula:



wherein P is the same as defined above;

- b. enolating the reaction product of step (a) produce a compound having the following formula:



wherein P and R are the same as defined above; and

05-31-06;01:40PM;

MAY 31 2006 ; 1-732-321-3030

# 5/ 12


IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/736,084	12/15/2003	Joseph C. Walsh	2003P88073US

Response To **OFFICIAL ACTION**

EXAMINER Krishnan, Ganapathy	
ART UNIT	PAGE NUMBER
1623	3

c. incorporating a leaving group to produce a compound having the following formula:



2. (Original) The method according to Claim 1, wherein P is selected from the group consisting of methoxymethyl ether, methylthiomethyl ether, 2-methoxyethoxymethyl ether, 1-ethoxyethyl ether, 1-methyl-1-methoxyethyl ether, t-butyl ether, allyl ether, benzyl ether, 4-nitrobenzyl ether, o-nitrobenzyl ether, trityl ether, monomethoxytrityl ether, dimethoxytrityl ether, tritylone ether, tetrahydropyran ether, tetrahydrothiopyranyl ether, 4-methoxy tetrahydropyran ether, 4-methoxytetrahydrothiopyranyl ether, tetrahydrofuran ether, tetrahydrotriofuranyl ether, isobutyrate ester, pivaloate ester, adamantate ester, benzoate ester, 2,4,6-trimethylbenzoate ester, methyl carbonate, allyl carbonate, benzyl carbonate, p-nitrobenzyl carbonate, t-Bu carbonate, S-benzylthio carbonate, N-phenyl carbamate, and nitrate ester.
3. (Original) The method according to Claim 1, wherein P is selected from the group consisting of dimethoxytrityl, monomethoxytrityl, trityl, t-butyloxycarbonyl, t-butyldimethylsilyl, t-butyldiphenylsilyl, tetrahydropyranyl ether, tetrahydrofuranyl ether, ethoxyethyl ether, and 1-methyl-1-methoxyethyl ether.
4. (Original) The method according to Claim 1, wherein R is alkyl C<sub>1</sub>-C<sub>4</sub>, i-propyl, benzyl, cycloalkane C<sub>3</sub>-C<sub>6</sub>, phenyl, tosyl, acetate, or benzoate.
5. (Original) The method according to Claim 1, wherein R is methyl, ethyl, i-propyl, benzyl, or cycloalkane C<sub>3</sub>-C<sub>6</sub>.

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/736,084	12/15/2003	Joseph C. Walsh	2003P88073US

## Response To OFFICIAL ACTION

EXAMINER	
Krishnan, Ganapathy	
ART UNIT	PAGE NUMBER
1623	4

6. (Original) The method according to Claim 1, wherein step (b) includes treating the reaction product of step (a) with an alkoxide having 1 to 4 carbons, cycloalkoxide C<sub>3</sub>-C<sub>6</sub>, phenoxide, tosylate, acetate, or benzoate.

7. (Original) The method according to Claim 6, wherein the alkoxide is sodium methoxide.

8. (Original) The method according to Claim 1, wherein L is a sulfonate ester.

9. (Original) The method according to Claim 1, wherein L is selected from the group consisting of mesylate, nosylate, tosylate, and triflate.

10. (Original) A method for preparing a precursor for the preparation of a radiolabeled nucleoside comprising:

- a. converting a 2-deoxy nucleoside into a 2,3'-anhydronucleoside;
- b. reacting the 2,3'-anhydronucleoside with a hydroxyl protecting group to produce a 2,3'-anhydronucleoside derivative wherein the 5'-O group is protected;
- c. reacting the protected 2,3'-anhydronucleoside derivative with a reagent that opens the 2,3'-anhydro-ring and enolates the 2-position on the pyrimidine ring; and
- d. incorporating a leaving group to produce the radiolabeled nucleoside precursor.

11. (Original) The method according to Claim 10, wherein the nucleoside is thymidine, cytidine, or uridine.

12. (Original) A method for preparing a precursor for the preparation of <sup>18</sup>F-FLT comprising:

- a. converting thymidine into 2,3'-anhydrothymidine;

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/736,084	12/15/2003	Joseph C. Walsh	2003P88073US

## Response To OFFICIAL ACTION

EXAMINER	
Krishnan, Ganapathy	
ART UNIT	PAGE NUMBER
1623	5

b. reacting the 2,3'-anhydro thymidine with a hydroxyl protecting group to produce a 2,3'-anhydrothymidine derivative wherein the 5'-O group is protected;

c. reacting the protected 2,3'-anhydrothymidine derivative with a reagent that opens the 2,3'-anhydro-ring and enolates the 2-position on the pyrimidine ring; and

d. incorporating a leaving group to produce the  $^{18}\text{F}$ -FLT precursor.

13. (Original) The method according to Claim 12, wherein step (c) produces an enol having an  $-\text{O}-\text{R}$  group attached to the 2-carbon.

14. (Original) A method according to Claim 13, wherein R is alkyl  $\text{C}_1-\text{C}_4$ , *i*-propyl, benzyl, cycloalkane  $\text{C}_3-\text{C}_6$ , phenyl, tosyl, acetate, or benzoate.

15. (Original) A method according to Claim 12, wherein step (c) includes treating the reaction product of step (b) with an alkoxide.

16. (Currently amended) A method according to Claim ~~15~~16, wherein the alkoxide is selected from the group consisting of sodium methoxide, and sodium ethoxide,

17. (Original) A method according to Claim 12, wherein the hydroxyl protecting group is dimethoxytrityl, monomethoxytrityl, trityl, *t*-butyloxycarbonyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl, tetrahydropyranyl ether, tetrahydrofuranyl ether, ethoxyethyl ether, or 1-methyl-1-methoxyethyl ether.

18. (Original) A method according to Claim 12, wherein the hydroxyl protecting group is dimethoxytrityl, monomethoxytrityl, or trityl.

19. (Original) A method according to Claim 12 wherein the leaving group is a sulfonate ester:

20. (Original) A method according to Claim 19, wherein the leaving group is mesylate, tosylate, nosylate, or triflate.


## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/736,084	12/15/2003	Joseph C. Walsh	2003P88073US

## Response To OFFICIAL ACTION

EXAMINER	
Krishnan, Ganapathy	
ART UNIT	PAGE NUMBER
1623	6

21. (Original) A compound having the following formula:



wherein R is alkyl C<sub>1</sub>-C<sub>4</sub>, *i*-propyl, benzyl, cycloalkane C<sub>3</sub>-C<sub>6</sub>, phenyl, tosyl, acetate, or benzoate; P is a hydroxyl protecting group; and L is a leaving group.

22. (Original) A compound according to Claim 21, wherein R is methyl or ethyl.
23. (Original) A compound according to Claim 21, wherein P is methoxymethyl ether, methylthiomethyl ether, 2-methoxyethoxymethyl ether, 1-ethoxyethyl ether, 1-methyl-1-methoxyethyl ether, *t*-butyl ether, allyl ether, benzyl ether, 4-nitrobenzyl ether, *o*-nitrobenzyl ether, trityl ether, monomethoxytrityl ether, dimethoxytrityl ether, tritylone ether, tetrahydropyran ether, tetrahydrothiopyranyl ether, 4-methoxy tetrahydropyran ether, 4-methoxytetrahydrothiopyranyl ether, tetrahydrofuran ether, tetrahydrotriofuranyl ether, isobutyrate ester, pivaloate ester, adamantate ester, benzoate ester, 2,4,6-trimethylbenzoate ester; methyl carbonate, allyl carbonate, benzyl carbonate, *p*-nitrobenzyl carbonate, *t*-Bu carbonate, *S*-benzylthio carbonate, *N*-phenyl carbamate, or nitrate ester.
24. (Original) A compound according to Claim 21, wherein P is dimethoxytrityl, monomethoxytrityl, trityl, *t*-butyloxycarbonyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl, tetrahydropyranyl ether, tetrahydrofuranyl ether, ethoxyethyl ether, or 1-methyl-1-methoxyethyl ether.
25. (Original) A compound according to Claim 21, wherein P is dimethoxytrityl.
26. (Original) A compound according to Claim 21, wherein L is a sulfonate ester.

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/736,084	12/15/2003	Joseph C. Walsh	2003P88073US

Response To **OFFICIAL ACTION**

EXAMINER
----------

Krishnan, Ganapathy

ART UNIT	PAGE NUMBER
----------	-------------


1623

7

27. (Original) A compound according to Claim 21, wherein L is selected from the group consisting of p-(2,4-dinitroanilino)benzenesulfonyl, benzenesulfonyl, methylsulfonyl (mesylate), p-methylbenzenesulfonyl (tosylate), 4-nitrobenzene sulfonyl (nosylate), p-bromobenzenesulfonyl, trifluoromethylsulfonyl (triflate), trichloroacetimidate, acyloxy, 2,2,2-trifluoroethanesulfonyl, imidazolesulfonyl, and 2,4,6-trichlorophenyl.


28. (Original) A compound according to Claim 21, wherein R is methyl, P is dimethoxy trityl, and L is mesylate, tosylate, or nosylate.

29. (Original) A compound having the following formula:



wherein Ms is methylsulfonyl.

30. (Original) A compound having the following formula:



wherein R is alkyl C<sub>1</sub>-C<sub>4</sub>, *i*-propyl, benzyl, cycloalkane C<sub>3</sub>-C<sub>6</sub>, phenyl, tosyl, acetate, or benzoate; P is a hydroxyl protecting group; X is oxygen, sulfur, or nitrogen, and L is a leaving group.

31. (Original) A compound according to Claim 30, wherein L is halogen, p-(2,4-dinitroanilino)benzenesulfonyl, benzenesulfonyl, methylsulfonyl (mesylate), p-methylbenzenesulfonyl (tosylate), 4-nitrobenzene sulfonyl (nosylate), p-

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.
10/736,084	12/15/2003	Joseph C. Walsh	2003P88073US

## Response To OFFICIAL ACTION

EXAMINER	
Krishnan, Ganapathy	
ART UNIT	PAGE NUMBER
1623	8

bromobenzenesulfonyl, trifluoromethylsulfonyl (triflate), trichloroacetimidate, acyloxy, 2,2,2-trifluoroethanesulfonyl, imidazolesulfonyl, or 2,4,6-trichlorophenyl.

32. (Original) A compound according to Claim 30, wherein P is dimethoxytrityl, monomethoxytrityl, trityl, t-butyloxycarbonyl, t-butyldimethylsilyl, t-butyldiphenylsilyl, tetrahydropyranyl ether, tetrahydrofuranyl ether, ethoxyethyl ether, or 1-methyl-1-methoxyethyl ether.

33-34. (Canceled)